


Granulomatous Meningoencephalomyelitis

(Inflammation of the Brain, Spinal Cord, and Meninges)

Basics

OVERVIEW

- “Granulomatous” refers to inflammation characterized by the presence of nodules; “meningoencephalomyelitis” is inflammation of the brain, spinal cord and their surrounding membranes (the membranes are known as “meninges”)
- GME was the most accepted and recognized central nervous system inflammatory disorder in the dog; accepted new terminology is Meningoencephalomyelitis of unknown origin (MUO)—many less serious viral and idiopathic disorders frequently were diagnosed erroneously as being “granulomatous meningoencephalomyelitis” (“idiopathic” means the disease is of unknown cause) but now are under the umbrella of the new classification, MUO
- MUO refers to an inflammatory disease that affects the central nervous system; it can be localized, widespread, or involve multiple locations
- Confirmation of the disease diagnosis is only possible through microscopic analysis of the affected nervous tissue, obtained through biopsy
- The spine is composed of multiple bones with disks (intervertebral disks) located in between adjacent bones (vertebrae); the disks act as shock absorbers and allow movement of the spine; the vertebrae are named according to their location—cervical vertebrae are located in the neck and are numbered as cervical vertebrae one through seven or C1–C7; thoracic vertebrae are located from the area of the shoulders to the end of the ribs and are numbered as thoracic vertebrae one through thirteen or T1–T13; lumbar vertebrae start at the end of the ribs and continue to the pelvis and are numbered as lumbar vertebrae one through seven or L1–L7; the remaining vertebrae are the sacral and coccygeal (tail) vertebrae



GENETICS

- A genetic basis is not proven

SIGNALMENT/DESCRIPTION OF PET

Species

- Dogs

Breed Predilections

- Historically, granulomatous meningoencephalomyelitis (now MUO) had been considered a disease of young to

middle-age, toy-breed dogs (especially terriers and poodles); however, any medium and larger breed can be affected

Mean Age and Range

- Mean—5 years of age
- Range, 6 months–10 years

Predominant Sex

- Females are slightly more likely to have this condition than are males

SIGNS/OBSERVED CHANGES IN THE PET

- Depend on the form of the disease and location of nerve tissue involved
- The cerebral form of disease (involving the brain) frequently results in seizure activity

Ocular Form

- Sudden (acute) onset of blindness, with dilated and unresponsive pupils; the “pupil” is the circular or elliptical opening in the center of the iris of the eye; the “iris” is the colored or pigmented part of the eye

Focal Form

- Cerebral lesions (involving the brain)—disorientation, behavioral changes, seizures, blindness, compulsive circling, head pressing
- Brainstem lesion (involving the part of the brain that is connected to the spinal cord that controls functions like breathing and heart rate)—drowsiness or sleepiness (known as “somnolence”), cranial nerve deficits (most commonly facial nerve deficit and lack of control of equilibrium, balance, and orientation [known as “vestibular dysfunction”]), one-sided weakness or partial paralysis (known as “hemiparesis”)
- Spinal cord lesion—neck pain, weakness or partial paralysis of all four legs (known as “tetraparesis”) for lesions in the spinal cord involving the area of the first cervical vertebra through the fifth cervical vertebra (C1–C5) or the sixth cervical vertebra through the second thoracic vertebra (C6–T2) or weakness or partial paralysis of the rear legs (known as “paraparesis”) for lesions in the spinal cord involving the area of the third thoracic vertebra through the third lumbar vertebra (T3–L3) or the fourth lumbar vertebra through the second sacral vertebra (L4–S2) and a wobbly, incoordinated, or “drunken”-appearing gait or movement (known as “ataxia”)

CAUSES

- Unknown

RISK FACTORS

- Unknown
- Some dogs develop clinical signs within 5–10 days of receiving vaccinations

Treatment

HEALTH CARE

- Stable pets can be discharged with recommended treatment
- Inpatient—for severely affected dogs; monitor pet closely for progression of nervous system deficits
- Intravenous fluids for pets that have lack of appetite (known as “anorexia”); fluids will be administered carefully to avoid over-hydration and worsening of fluid buildup in the brain (known as “cerebral edema”)
- Provide a padded cage for dogs with lack of control of equilibrium, balance, and orientation (vestibular dysfunction), severe dementia or seizure activity
- Recumbent pets should be turned frequently (every 4 hours) to avoid pressure sores and lung congestion

ACTIVITY

- Depends on severity of disease and location of nerve tissue involved
- Pets with a wobbly, incoordinated or “drunken” appearing gait or movement (ataxia) should be confined to a padded cage to avoid injury

DIET

- Ensure adequate caloric intake

Medications

Medications presented in this section are intended to provide general information about possible treatment. The treatment for a particular condition may evolve as medical advances are made; therefore, the medications should not be considered as all inclusive

- Steroids—dexamethasone followed by prednisone; the steroid dose should be adjusted by your pet's veterinarian, according to response to treatment and side effects—the goal is to find the dose that keeps the clinical signs controlled with minimal side effects
- To prevent gastrointestinal ulceration, administer omeprazole (proton pump inhibitor) and famotidine (an H₂-blocker) to reduce stomach acid with the steroid therapy
- Phenobarbital, zonisamide or levetiracetam to control seizures
- Gabapentin for compulsive circling
- Chemotherapeutic drugs used to decrease the immune response—azathioprine, cytosine arabinoside
- Medications to decrease the immune response (known as “immunosuppressive drugs”)—cyclosporine, leflunomide
- Radiation therapy is an alternative treatment in the focal form of the disease when other therapies have failed (the diagnosis should be confirmed by microscopic evaluation of tissue samples obtained by biopsy, before starting radiation therapy)

Follow-Up Care

PATIENT MONITORING

- Repeat nervous system examination periodically (every 2–4 weeks)
- Evaluate bloodwork (complete blood count [CBC] and biochemical profile) regularly to monitor for low white blood cell count (known as “leukopenia”), low platelet count (known as “thrombocytopenia”), and liver and kidney function, if chemotherapy or medications to decrease the immune response (immunosuppressive drugs) are included in treatment
- Monitor urine in pets on long-term steroid treatment—protein in the urine (known as “proteinuria”) or urinary tract infection are frequent consequences of long-term steroid treatment

POSSIBLE COMPLICATIONS

- Deterioration of clinical signs, despite aggressive treatment
- Repeated or prolonged seizure activity (known as “status epilepticus”)
- Dementia
- Brain pushes downward in the skull and herniates through the opening that leads to the neck (known as “tentorial herniation” or “brain herniation”) and death

EXPECTED COURSE AND PROGNOSIS

- Not all pets with central nervous system inflammatory disease have a poor prognosis
- Granulomatous meningoencephalomyelitis (now MUO) has been characterized as a fatal disease without enough scientific evidence; it is uncertain if dogs that survive inflammatory central nervous system disease had GME, as brain biopsies rarely are obtained to provide a definitive diagnosis; about 70% are expected to respond to therapy, especially if started early in the course of the disease

Key Points

- Clinical signs overlap significantly among different central nervous system inflammatory diseases—a diagnostic workup is very important
- Mortality rate for MUO is clearly biased by the severe cases that go to post-mortem examinations; brain biopsies rarely are obtained during life to provide a definitive diagnosis and to provide a more accurate mortality rate
- Some dogs with central nervous system inflammatory disease can be treated successfully; however, long-term treatment and client compliance are required
- Steroid therapy may be necessary indefinitely